

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID: SSSPTA1626GMS

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

NEWS 1		Web Page for STN Seminar Schedule - N. America
NEWS 2	AUG 10	Time limit for inactive STN sessions doubles to 40 minutes
NEWS 3	AUG 18	COMPENDEX indexing changed for the Corporate Source (CS) field
NEWS 4	AUG 24	ENCOMPPLIT/ENCOMPPLIT2 reloaded and enhanced
NEWS 5	AUG 24	CA/Caplus enhanced with legal status information for U.S. patents
NEWS 6	SEP 09	50 Millionth Unique Chemical Substance Recorded in CAS REGISTRY
NEWS 7	SEP 11	WPIDS, WPINDEX, and WPIX now include Japanese FTERM thesaurus
NEWS 8	OCT 21	Derwent World Patents Index Coverage of Indian and Taiwanese Content Expanded
NEWS 9	OCT 21	Derwent World Patents Index enhanced with human translated claims for Chinese Applications and Utility Models
NEWS 10	NOV 23	Addition of SCAN format to selected STN databases
NEWS 11	NOV 23	Annual Reload of IFI Databases
NEWS 12	DEC 01	FRFULL Content and Search Enhancements
NEWS 13	DEC 01	DGENE, USGENE, and PCTGEN: new percent identity feature for sorting BLAST answer sets
NEWS 14	DEC 02	Derwent World Patent Index: Japanese FI-TERM thesaurus added
NEWS 15	DEC 02	PCTGEN enhanced with patent family and legal status display data from INPADOCDB
NEWS 16	DEC 02	USGENE: Enhanced coverage of bibliographic and sequence information
NEWS 17	DEC 21	New Indicator Identifies Multiple Basic Patent Records Containing Equivalent Chemical Indexing in CA/Caplus

NEWS EXPRESS MAY 26 09 CURRENT WINDOWS VERSION IS V8.4,
AND CURRENT DISCOVER FILE IS DATED 06 APRIL 2009

NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS LOGIN Welcome Banner and News Items

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN customer agreement. This agreement limits use to scientific research. Use for software development or design, implementation of commercial gateways, or use of CAS and STN data in the building of commercial products is prohibited and may result in loss of user privileges and other penalties.

FILE 'HOME' ENTERED AT 15:20:36 ON 04 JAN 2010

\Rightarrow

Uploading

THIS COMMAND NOT AVAILABLE IN THE CURRENT F

Do you want to

Choice (Y/n):

Some commands only work in certain files. For example, the EXPAND command can only be used to look at the index in a file which has an index. Enter "HELP COMMANDS" at an arrow prompt (=>) for a list of commands which can be used in this file.

→ FILE REGISTRY

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	0.22	0.22

FILE 'REGISTRY' ENTERED AT 15:21:02 ON 04 JAN 2010
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2010 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 3 JAN 2010 HIGHEST RN 1200115-43-0
DICTIONARY FILE UPDATES: 3 JAN 2010 HIGHEST RN 1200115-43-0

New CAS Information Use Policies, enter **HELP USAGETERMS** for details.

TSCA INFORMATION NOW CURRENT THROUGH June 26, 2009.

Please note that search-term pricing does apply when conducting SmartSELECT searches.

REGISTRY includes numerically searchable data for experimental and predicted properties as well as tags indicating availability of experimental property data in the original document. For information on property searching in REGISTRY, refer to:

<http://www.cas.org/support/stnqgen/stndoc/properties.html>

=>
Uploading C:\Program Files\Stnexp\Queries\10591921a.str



chain nodes :
17 18 19 20 21 22 23 24 26 27 29 35 36

10591921

ring nodes :
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 25 30 31 32 33 34
chain bonds :
1-17 3-21 4-19 4-20 5-18 6-22 7-23 8-27 12-26 15-24 24-29 24-36 29-30
34-35
ring bonds :
1-2 1-16 2-3 3-4 4-5 5-6 6-7 7-8 8-9 9-10 10-11 11-12 12-13 12-25
13-14 13-25 14-15 15-16 30-31 30-33 31-32 32-34 33-34
exact/norm bonds :
1-2 1-16 1-17 2-3 3-4 3-21 4-5 4-19 4-20 5-6 5-18 6-7 6-22 7-8 7-23
8-9 8-27 9-10 10-11 11-12 12-13 12-25 12-26 13-14 13-25 14-15 15-16
15-24 24-29 24-36 29-30 30-31 30-33 31-32 32-34 33-34 34-35
isolated ring systems :
containing 1 : 30 :

G1:C,O

G2:H,Ak,CH3,Et,n-Pr

Match level :

1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:Atom 8:Atom 9:Atom 10:Atom
11:Atom 12:Atom 13:Atom 14:Atom 15:Atom 16:Atom 17:CLASS 18:CLASS 19:CLASS
20:CLASS 21:CLASS 22:CLASS 23:CLASS 24:CLASS 25:Atom 26:CLASS 27:CLASS
29:CLASS 30:Atom 31:Atom 32:Atom 33:Atom 34:Atom 35:CLASS 36:CLASS

Stereo Bonds:

22-6 (Single Hash).
23-7 (Single Wedge).
24-15 (Single Wedge).
27-8 (Single Hash).

Stereo Chiral Centers:

6 (Parity=Even)
7 (Parity=Odd)
8 (Parity=Odd)
15 (Parity=Odd)

Stereo RSS Sets:

Type=Relative (Default). 4 Nodes= 6 7 8 15

L1 STRUCTURE UPLOADED

=> d 11
L1 HAS NO ANSWERS
L1 STR
*** STRUCTURE DIAGRAM IS NOT AVAILABLE ***

Structure attributes must be viewed using STN Express query preparation.

=> s 11
SAMPLE SEARCH INITIATED 15:21:31 FILE 'REGISTRY'

10591921

SAMPLE SCREEN SEARCH COMPLETED - 45 TO ITERATE
100.0% PROCESSED 45 ITERATIONS 10 ANSWERS
SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS: ONLINE **COMPLETE**
BATCH **COMPLETE**
PROJECTED ITERATIONS: 498 TO 1302
PROJECTED ANSWERS: 11 TO 389

L2 10 SEA SSS SAM L1

=> s 11 sss full
FULL SEARCH INITIATED 15:21:37 FILE 'REGISTRY'
FULL SCREEN SEARCH COMPLETED - 770 TO ITERATE

100.0% PROCESSED 770 ITERATIONS 175 ANSWERS
SEARCH TIME: 00.00.01

L3 175 SEA SSS FUL L1

=> FIL HCAPLUS
COST IN U.S. DOLLARS SINCE FILE TOTAL
FULL ESTIMATED COST ENTRY SESSION
191.54 191.76

FILE 'HCAPLUS' ENTERED AT 15:21:42 ON 04 JAN 2010
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2010 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 4 Jan 2010 VOL 152 ISS 2
FILE LAST UPDATED: 3 Jan 2010 (20100103/ED)
REVISED CLASS FIELDS (/NCL) LAST RELOADED: Oct 2009
USPTO MANUAL OF CLASSIFICATIONS THESAURUS ISSUE DATE: Oct 2009

HCplus now includes complete International Patent Classification (IPC) reclassification data for the third quarter of 2009.

CAS Information Use Policies apply and are available at:

<http://www.cas.org/legal/infopolicy.html>

This file contains CAS Registry Numbers for easy and accurate substance identification.

10591921

=> s 13
L4 832 L3

=> s 14 and epothilone derivatives
1258 EPOTHILONE
1020 EPOTHILONES
1626 EPOTHILONE
(EPOTHILONE OR EPOTHILONES)
379022 DERIVATIVES
1242341 DERIVS
1359744 DERIVATIVES
(DERIVATIVES OR DERIVS)
106 EPOTHILONE DERIVATIVES
(EPOTHILONE(W) DERIVATIVES)
L5 71 L4 AND EPOTHILONE DERIVATIVES

=> s 15 and py<=2004
25162081 PY<=2004
L6 48 L5 AND PY<=2004

=> s 16 and p/dt
7012674 P/DT
L7 45 L6 AND P/DT

=> s 17 and us/pc
2024782 US/PC
L8 38 L7 AND US/PC

=> s 18 and pharmaceutical
376608 PHARMACEUTICAL
95011 PHARMACEUTICALS
433406 PHARMACEUTICAL
(PHARMACEUTICAL OR PHARMACEUTICALS)
L9 16 L8 AND PHARMACEUTICAL

=> d 19 ibib abs hitstr tot

L9 ANSWER 1 OF 16 HCAPLUS COPYRIGHT 2010 ACS on STN
ACCESSION NUMBER: 2005:572598 HCAPLUS
DOCUMENT NUMBER: 143:97209
TITLE: Synthesis of epothilones for use in pharmaceutical compositions as antitumor agents
INVENTOR(S): Danishefsky, Samuel J.; Rivkin, Alexey; Yoshimura, Fumihiro; Chou, Ting-Chao; Gabarda, Ana E.; Dong, Huajin; Wu, Kaida; Moore, Malcolm A. S.; Dorn, David USA
PATENT ASSIGNEE(S): U.S. Pat. Appl. Publ., 274 pp., Cont.-in-part of U.S. Ser. No. 435,408.
SOURCE: CODEN: USXXCO
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 3
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	-----

US 20050143429	A1	20050630	US 2004-921109	20040818 <--
US 7384964	B2	20080610		
US 20040053995	A1	20040318	US 2003-402004	20030328 <--
US 6921769	B2	20050726		
US 20040053910	A1	20040318	US 2003-435408	20030509 <--
AU 2005218308	A1	20050915	AU 2005-218308	20050228
CA 2556692	A1	20050915	CA 2005-2556692	20050228
WO 2005084222	A2	20050915	WO 2005-US6051	20050228
WO 2005084222	A3	20051124		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
EP 1722791	A2	20061122	EP 2005-723768	20050228
R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR				
CN 1976699	A	20070606	CN 2005-80013480	20050228
BR 2005008232	A	20070717	BR 2005-8232	20050228
JP 2007525519	T	20070906	JP 2007-500999	20050228
ZA 2005002337	A	20060927	ZA 2005-2337	20050318
MX 2006009792	A	20061116	MX 2006-9792	20060828
KR 2007100626	A	20071011	KR 2006-720063	20060927
US 20090149516	A1	20090611	US 2008-135823	20080609 <--
PRIORITY APPLN. INFO.:				
US 2002-405823P P 20020823				
US 2002-408589P P 20020906				
US 2002-423129P P 20021101				
US 2003-456159P P 20030320				
US 2003-402004 A2 20030328				
US 2003-435408 A2 20030509				
US 2003-496741P P 20030821				
US 2004-548402P P 20040227				
US 2004-921109 A 20040818				
WO 2005-US6051 W 20050228				

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OTHER SOURCE(S): CASREACT 143:97209; MARPAT 143:97209

GI



AB Epothilone analogs, such as I [-A-B-, -C-D- = -C.tplbond.C-, -CH(R)CH(R1)-, -(C(R)C(R1))-; R, R1 = H, alkyl, halogen, alkoxy, acyl, etc.; -A-B- = fused oxirane ring; -C-D- = fused cyclopropane or fused aziridine ring; R2 = aryl, heteroaryl, arylalkyl, heteroarylkylalkyl] are prepared as antitumor agents. The present invention also provides pharmaceutical compns. comprising compds. of formula I and provides methods of treating cancer comprising administering a compound of formula I. Thus, II was prepared via an intramol. methathesis macrocyclization synthetic sequence and showed good cell growth inhibition against various drug-resistant tumors.

IT 152044-54-7P 190370-13-9P


RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(synthesis of epothilone derivs. for use in pharmaceutical compns. as antitumor agents)

RN 152044-54-7 HCAPLUS

CN 4,17-Dioxabicyclo[14.1.0]heptadecane-5,9-dione,
7,11-dihydroxy-8,8,10,12,16-pentamethyl-3-[(1E)-1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]- (1S,3S,7S,10R,11S,12S,16R)- (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).
Double bond geometry as shown.




RN 190370-13-9 HCPLUS

CN 4,17-Dioxabicyclo[14.1.0]heptadecane-5,9-dione,
7,11-dihydroxy-8,10,12,16-pentamethyl-3-[(1E)-1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-, (1R,3S,7S,10R,11S,12S,16S)- (CA INDEX NAME)

Absolute stereochemistry.

Double bond geometry as shown.




REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L9 ANSWER 2 OF 16 HCPLUS COPYRIGHT 2010 ACS on STN
 ACCESSION NUMBER: 2004:780536 HCPLUS
 DOCUMENT NUMBER: 141:271549
 TITLE: Treatment of proliferative diseases with
 epothilone derivatives and radiation
 INVENTOR(S): Fruschy, Martin
 PATENT ASSIGNEE(S): Novartis A.-G., Switz.; Novartis Pharma G.m.b.H.
 SOURCE: PCT Int. Appl., 17 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004080458	A1	20040923	WO 2004-EP2610	20040312 <--
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
CA 2519037	A1	20040323	CA 2004-2519037	20040312 <--
AU 2004218927	A1	20040923	AU 2004-218927	20040312 <--
EP 1605937	A1	20051221	EP 2004-719972	20040312
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK				
BR 2004008366	A	20060321	BR 2004-8366	20040312
CN 1758908	A	20060412	CN 2004-80006581	20040312
JP 2006520353	T	20060907	JP 2006-504672	20040312
NZ 542305	A	20090331	NZ 2004-542305	20040312
ZA 2005006942	A	20060726	ZA 2005-6942	20050830
NO 2005004723	A	20051013	NO 2005-4723	20051013
US 20070129411	A1	20070607	US 2006-549978	20061115 <--
AU 2008200555	A1	20080228	AU 2008-200555	20080206
PRIORITY APPLN. INFO.:			GB 2003-5928	A 20030314
			AU 2004-218927	A3 20040312
			WO 2004-EP2610	W 20040312

OTHER SOURCE(S): MARPAT 141:271549
 GI



I

AB The invention discloses compds. I (A = O, NRN [RN = H, lower alkyl]; R = H, lower alkyl; Z = O, bond], in particular pharmaceutical compns. for use in combination with ionizing radiation for the delay of progression or treatment of a proliferative disease, especially a solid tumor disease.

IT 152044-54-7, Epothilone B


RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(Treatment of proliferative diseases with epothilone derivs. and radiation)

RN 152044-54-7 HCAPLUS

CN 4,17-Dioxabicyclo[14.1.0]heptadecane-5,9-dione, 7,11-dihydroxy-8,8,10,12,16-pentamethyl-3-[(1E)-1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-, (1S,3S,7S,10R,11S,12S,16R)- (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).
Double bond geometry as shown.



OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD
(1 CITINGS)

REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L9 ANSWER 3 OF 16 HCPLUS COPYRIGHT 2010 ACS on STN
 ACCESSION NUMBER: 2004:550960 HCPLUS
 DOCUMENT NUMBER: 141:106321
 TITLE: Preparation of epothilone
 derivatives for use in pharmaceutical
 compositions as antitumor agents
 INVENTOR(S): Denni-Dischert, Donatiinne; Floersheimer, Andreas;
 Kuesters, Ernst; Oberer, Lukas; Sedelmeier, Gottfried
 PATENT ASSIGNEE(S): Novartis A.-G., Switz.; Novartis Pharma G.m.b.H.
 SOURCE: PCT Int. Appl., 50 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004056832	A2	20040708	WO 2003-EP14747	20031222 <--
WO 2004056832	A3	20040910		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, HR, HU, ID, IL, IN, IS, JP, KE, KG, KE, KR, KZ, LC, LK, LT, LU, LV, MA, MD, MK, MN, MX, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SE, SG, SK, SY, TJ, TM, TN, TR, TT, UA, US, UZ, VC, VN, YU, ZA, ZW				
RW: AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR				
CA 2510620	A1	20040708	CA 2003-2510620	20031222 <--
AU 2003294938	A1	20040714	AU 2003-294938	20031222 <--
EP 1581536	A2	20051005	EP 2003-785920	20031222
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
BR 2003017693	A	20051122	BR 2003-17693	20031222
CN 1732172	A	20060208	CN 2003-80107416	20031222
CN 100384851	C	20080430		
JP 2006514025	T	20060427	JP 2004-561416	20031222
US 20060014796	A1	20060119	US 2005-538200	20050609 <--
US 7317100	B2	20080108		
PRIORITY APPLN. INFO.:			GB 2002-30024	A 20021223
			WO 2003-EP14747	W 20031222

OTHER SOURCE(S): MARPAT 141:106321
 GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB C4-demethyl-epothilones or C4-bisnor-epothilones, such as I [R1, R7 = H, alkyl; R2 = nitrogen containing heteroaryl; R3 = H, Me; X = O, NR7; Z = O, bond], were prepared via fermentation and organic synthesis for use in pharmaceutical compns. as antitumor agents. Thus,

C4-bisnor-epothilone B II ($R_3 = H$) was prepared via an aldol condensation of aldehyde III with *in situ* disilylated (3*R*)-3-hydroxy-5-oxoheptanoic acid followed by a desilylation/macrolactonization reaction sequence. Also, C4-demethyl-epothilone B II ($R = Me$) was prepared directly by a fermentation process. The prepared epothilones were assayed for efficacy against human KB-31 and KB-8511 carcinoma cells. Drug delivery formulations containing the prepared epothilones were presented.

IT 152044-53-6, Epothilone A 152044-54-7, Epothilone B


RL: PAC (Pharmacological activity); RCT (Reactant); THU (Therapeutic use); BIOL (Biological study); RACT (Reactant or reagent); USES (Uses) (preparation of epothilone derivs. via fermentation and organic synthesis for use in pharmaceutical compns. as antitumor agents)

RN 152044-53-6 HCPLUS

CN 4,17-Dioxabicyclo[4.1.0]heptadecane-5,9-dione, 7,11-dihydroxy-8,8,10,12-tetramethyl-3-[(1*E*)-1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-, (1*S*,3*S*,7*S*,10*R*,11*S*,12*S*,16*R*)- (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).

Double bond geometry as shown.




RN 152044-54-7 HCPLUS

CN 4,17-Dioxabicyclo[4.1.0]heptadecane-5,9-dione, 7,11-dihydroxy-8,8,10,12,16-pentamethyl-3-[(1*E*)-1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-, (1*S*,3*S*,7*S*,10*R*,11*S*,12*S*,16*R*)- (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).

Double bond geometry as shown.



REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L9 ANSWER 4 OF 16 HCPLUS COPYRIGHT 2010 ACS on STN
ACCESSION NUMBER: 2004:428797 HCPLUS

DOCUMENT NUMBER: 141:12271

TITLE: Preparation of protein-stabilized liposomal formulations of pharmaceutical agents

INVENTOR(S): Singh, Chandra U.

PATENT ASSIGNEE(S): Azaya Therapeutics, Inc., USA

SOURCE: PCT Int. Appl., 40 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004043363	A2	20040527	WO 2003-US35297	20031106 <--
WO 2004043363	A3	20040812		
W: AB, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
CA 2505520	A1	20040527	CA 2003-2505520	20031106 <--

AU 2003287526	A1 20040603	AU 2003-287526	20031106 <--
US 20040247660	A1 20041209	US 2003-703187	20031106 <--
US 7179484	B2 20070220		
EP 1585504	A2 20051019	EP 2003-781768	20031106
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK			
JP 2006508126	T 20060309	JP 2004-551766	20031106
US 20070166368	A1 20070719	US 2007-619526	20070103 <--
PRIORITY APPLN. INFO.:		US 2002-424230P	P 20021106
		US 2003-703187	A1 20031106
		WO 2003-US35297	W 20031106

AB The present invention provides a method of preparing a protein-stabilized lipid formulation containing at least one lipophilic pharmaceutical agent. Specifically, the present invention discloses compns. and methods for protein stabilized liposomes, the creation of protein stabilized liposomes, and the administration of protein stabilized liposomes.

IT 152044-53-6, Epothilone-A 152044-54-7, Epothilone-B


RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(preparation of protein-stabilized liposomal formulations of pharmaceutical agents)

RN 152044-53-6 HCPLUS

CN 4,17-Dioxabicyclo[14.1.0]heptadecane-5,9-dione,
7,11-dihydroxy-8,8,10,12-tetramethyl-3-[(1E)-1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-, (1S,3S,7S,10R,11S,12S,16R)- (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).


Double bond geometry as shown.



RN 152044-54-7 HCPLUS

CN 4,17-Dioxabicyclo[14.1.0]heptadecane-5,9-dione,
7,11-dihydroxy-8,8,10,12,16-pentamethyl-3-[(1E)-1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-, (1S,3S,7S,10R,11S,12S,16R)- (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).
Double bond geometry as shown.




OS.CITING REF COUNT: 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD
(2 CITINGS)
REFERENCE COUNT: 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L9 ANSWER 5 OF 16 HCPLUS COPYRIGHT 2010 ACS on STN
ACCESSION NUMBER: 2004:331975 HCPLUS
DOCUMENT NUMBER: 140:332484
TITLE: Epothilone derivatives for the
treatment of multiple myeloma
INVENTOR(S): Lin, Boris; Anderson, Kenneth C.; Griffin, James
Douglas
PATENT ASSIGNEE(S): Dana-Farber Cancer Institute Inc., USA
SOURCE: PCT Int. Appl., 23 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004032923	A1	20040422	WO 2003-IB4480	20031010 <--
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LT, LU, LV, MA, MD, MK, MN, MX, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SE, SG, SK, SY, TJ, TM, TN, TR, TT, UA, US, UZ, VC, VN, YU, ZA, ZW				
RW: AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE,				

DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR				
CA 2501610	A1	20040422	CA 2003-2501610	20031010 <--
AU 2003264822	A1	20040504	AU 2003-264822	20031010 <--
EP 1553939	A1	20050720	EP 2003-807949	20031010
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
BR 2003014555	A	20050809	BR 2003-14555	20031010
CN 1703216	A	20051130	CN 2003-80101065	20031010
JP 20060504727	T	20060209	JP 2004-542749	20031010
US 20060094766	A1	20060504	US 2005-530857	20050816 <--
US 20090233973	A1	20090917	US 2009-474555	20090529 <--
PRIORITY APPLN. INFO.:			US 2002-417916P	P 20021011
			WO 2003-IB4480	W 20031010
			US 2005-530857	A1 20050816

OTHER SOURCE(S): MARPAT 140:332484
GI



AB The invention discloses a method of treating a warm-blooded animal, especially a human, having myeloma, especially myeloma which is resistant to conventional cytotoxic chemotherapy, comprising administering to said animal a therapeutically effective amount of an epothilone, especially an epothilone of formula I (wherein A = O or NR, wherein RN = hydrogen or lower alkyl, R = hydrogen or lower alkyl, R' = Me, methoxy, ethoxy, amino, methylamino, dimethylamino or methylthio, and Z = O or bond), to a combination comprising an epothilone, for simultaneous, sep. or sequential use; and to a pharmaceutical composition and a com. package comprising said combination.


IT 152044-54-7, Epothilone B

RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(epothilone derivs. for treatment of multiple myeloma)

RN 152044-54-7 HCPLUS

CN 4,17-Dioxabicyclo[14.1.0]heptadecane-5,9-dione,
7,11-dihydroxy-8,8,10,12,16-pentamethyl-3-[(1E)-1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-, (1S,3S,7S,10R,11S,12S,16R)- (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).
Double bond geometry as shown.



OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD
(1 CITINGS)

REFERENCE COUNT: 9 THERE ARE 9 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L9 ANSWER 6 OF 16 HCPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2004:117139 HCPLUS
DOCUMENT NUMBER: 140:181250

TITLE: Preparation of new epothilone peptide effector conjugates for pharmaceutical use in the treatment of proliferative or angiogenesis associated disease processes

INVENTOR(S): Berger, Markus; Klar, Ulrich; Siemeister, Gerhard; Willuda, Joerg; Menrad, Andreas; Bosslet, Klaus

PATENT ASSIGNEE(S): Schering AG, Germany
SOURCE: Ger. Offen., 43 pp.

DOCUMENT TYPE: Patent
LANGUAGE: German

FAMILY ACC. NUM. COUNT: 2

PATENT INFORMATION:



PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 10234975	A1	20040212	DE 2002-10234975	20020731 <--
CA 2492437	A1	20040212	CA 2003-2492437	20030731 <--
WO 2004012735	A2	20040212	WO 2003-EP8483	20030731 <--
WO 2004012735	A3	20040527		
W:	AB, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG,			

PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR,
 TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW
 RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY,
 KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES,
 FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR,
 BE, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG
 AU 2003253365 A1 20040223 AU 2003-253365 20030731 <--
 US 20050026971 A1 20050203 US 2003-631011 20030731 <--
 US 7129254 B2 20061031
 EP 1524979 A2 20050427 EP 2003-743752 20030731
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LT, LU, NL, SE, MC, PT,
 IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK
 BR 2003013043 A 20050614 BR 2003-13043 20030731
 JP 2006505627 T 20060216 JP 2005-506073 20030731
 CN 1812785 A 20060802 CN 2003-818111 20030731
 NZ 537870 A 20070330 NZ 2003-537870 20030731
 IN 2005DN00038 A 20090320 IN 2005-DN38 20050105
 MX 2005001282 A 20050428 MX 2005-1282 20050131
 ZA 2005001648 A 20060531 ZA 2005-1648 20050224
 NO 2005001038 A 20050406 NO 2005-1038 20050225
 US 20070088060 A1 20070419 US 2006-509557 20060825 <--
 US 7335775 B2 20080226
 IN 2007DN04669 A 20070831 IN 2007-DN4669 20070618
 US 20080166362 A1 20080710 US 2007-258 20071211 <--
 PRIORITY APPLN. INFO.:
 DE 2002-10234975 A 20020731
 DE 2003-10305098 A 20030207
 US 2003-451673P P 20030305
 US 2003-631011 A3 20030731
 WO 2003-EP8483 W 20030731
 IN 2005-DN38 A3 20050105
 US 2006-509557 A3 20060825

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OTHER SOURCE(S): MARPAT 140:181250

GI



AB Preparation of epothilone derivs., such as I (R2 = alkyl, alkenyl, alkynyl, aryl, etc.; L1, L2 = carboxyl, carbamoyl, carbonic linking group with a terminal group, such as maleimido, suitable for forming a sulfide link with a bio. mol.; L3 = heteroaryl, such as thiazol-4-yl; W = alkenylene linking group; or L3W = heteroaryl, such as benzothiazol-5-yl; X = O, bond), as effectors linked with suitable biomols. as recognition units was described (no biol. testing data was presented). Production of the epothilone conjugates was carried out by the effectors being reacted with suitable linkers, and the compds. that were produced were conjugated to biomol. recognition units. These conjugates are claimed for use in the treatment of proliferative or angiogenesis-associated disease processes, such as tumors, inflammatory diseases, neurodegenerative diseases, such as multiple sclerosis and Alzheimer's disease, and rheumatoid arthritis. Thus, epothilone derivative II [L1 = 3-(2,5-dioxo-2,5-dihydropyrrol-1-yl)-1-Pr, L2 = H] was prepared via a carbamoylation of silylated epothilone I (L1 = H, L2 = SiMe2CMe3) with 3-(2,5-dioxo-2,5-dihydropyrrol-1-yl)-1-propylisocyanate and subsequent desilylation.


IT 152044-54-7

RL: RCT (Reactant); RACT (Reactant or reagent)
(preparation of new epothilone antibody peptide effector conjugates for pharmaceutical use in the treatment of proliferative or angiogenesis associated disease processes)

RN 152044-54-7 HCAPLUS


CN 4,17-Dioxabicyclo[4.1.0]heptadecane-5,9-dione,
7,11-dihydroxy-8,8,10,12,16-pentamethyl-3-[(1E)-1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-, (1S,3S,7S,10R,11S,12S,16R)- (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).
Double bond geometry as shown.



IT 152044-54-7DP, sulfide conjugate with peptide recognition biomol. 220773-47-7DP, sulfide conjugate with peptide recognition biomol.
 RL: SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of new epothilone antibody peptide effector conjugates for pharmaceutical use in the treatment of proliferative or angiogenesis associated disease processes)
 RN 152044-54-7 HCAPLUS
 CN 4,17-Dioxabicyclo[4.1.0]heptadecane-5,9-dione,
 7,11-dihydroxy-8,10,12,16-pentamethyl-1-[(1E)-1-methyl-2-(2-methyl-4-thiazolyl)ethyl]-, (1S,3S,7S,10R,11S,12S,16R)- (CA INDEX NAME)


Absolute stereochemistry. Rotation (-).
Double bond geometry as shown.



RN 220773-47-7 HCPLUS

CN 4,17-Dioxabicyclo[14.1.0]heptadecane-5,9-dione,
 10-ethyl-7,11-dihydroxy-8,8,12,16-tetramethyl-3-[(1E)-1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-, (1S,3S,7S,10R,11S,12S,16R)- (CA INDEX NAME)

Absolute stereochemistry.
 Double bond geometry as shown.




OS.CITING REF COUNT: 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD
(2 CITINGS)

L9 ANSWER 7 OF 16 HCPLUS COPYRIGHT 2010 ACS on STN
ACCESSION NUMBER: 2003:931114 HCPLUS
DOCUMENT NUMBER: 139:395751
TITLE: Preparation of C-21 modified epothilone derivatives for use in pharmaceutical compositions for the treatment of cancer
INVENTOR(S): Lee, Francis Y. F.; Haby, Thomas A.; Naringrekar, Vijay H.; Raghavan, Krishnaswamy S.; Franchini, Miriam K.
PATENT ASSIGNEE(S): Bristol-Myers Squibb Company, USA
SOURCE: PCT Int. Appl., 65 pp.
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003096975	A2	20031127	WO 2003-US15097	20030513 <--
WO 2003096975	A3	20031224		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
AU 2003234545	A1	20031202	AU 2003-234545	20030513 <--
US 20040053978	A1	20040318	US 2003-437103	20030513 <--
US 7053069	B2	20060530		
EP 1505969	A2	20050216	EP 2003-728885	20030513
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LT, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
PRIORITY APPLN. INFO.:			US 2002-380634P	P 20020515
			WO 2003-US15097	W 20030513

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OTHER SOURCE(S): MARPAT 139:395751
GI



AB C-21 modified epothilones, such as I [R = NH₂, OH, SH, alkylamino, alkoxy, alkylthio, etc.], were prepared for therapeutic use as antitumor agents. Thus, 21-aminoepothilone B I (R = NH₂) was prepared by reaction of epothilone F I (R = OH) with diphenylphosphoryl azide in THF under argon to give 21-azidoepothilone B I (R = N₃) in 91% yield and subsequent hydrogenation of the azide using Lindlar catalyst in EtOH and an H₂ atmosphere to give the target amine in 81% yield. The compns. are stable and readily prepared for administration by dissoln. in aqueous vehicles suitable for i.v. administration. A process for formulating C-21 modified epothilone derivs. for oral and parenteral administration was disclosed.

IT 152044-54-7, Epothilone B


RL: RCT (Reactant); RACT (Reactant or reagent)

(preparation of C-21 modified epothilone derivs. for use in pharmaceutical compns. for treatment of cancer)

RN 152044-54-7 HCAPLUS


CN 4,17-Dioxabicyclo[4.1.0]heptadecane-5,9-dione,
7,11-dihydroxy-8,10,12,16-pentamethyl-3-[(1E)-1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-, (1S,3S,7S,10R,11S,12S,16R)- (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).
Double bond geometry as shown.



IT 219990-27-9P, Epothilone B N-oxide
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT
 (Reactant or reagent)
 (preparation of C-21 modified epothilone derivs. for use
 in pharmaceutical compns. for treatment of cancer)
 RN 219990-27-9 HCPLUS
 CN 4,17-Dioxabicyclo[4.1.0]heptadecane-5,9-dione,
 7,11-dihydroxy-8,8,10,12,16-pentamethyl-3-[(1E)-1-methyl-2-(2-methyl-3-
 oxido-4-thiazolyl)ethenyl]-, (1S,3S,7S,10R,11S,12S,16R)- (CA INDEX NAME)

Absolute stereochemistry.
 Double bond geometry as shown.



OS.CITING REF COUNT: 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD
(2 CITINGS)

REFERENCE COUNT: 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L9 ANSWER 8 OF 16 HCPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2003:757689 HCPLUS

DOCUMENT NUMBER: 139:276755

TITLE: Preparation of epothilone derivatives for therapeutic use as anticancer agents

INVENTOR(S): Regueiro-Ren, Alicia; Kim, Soong-Hoon

PATENT ASSIGNEE(S): Bristol-Myers Squibb Company, USA

SOURCE: PCT Int. Appl., 47 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1


PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003078411	A1	20030925	WO 2003-US7584	20030311 <--
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR,				

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OTHER SOURCE(S): MARPAT 139:276755

GI



AB Epothilone derivs., such as I [M = bond, O, NR9, CR1OR11; X = O, NH; R1-R4 = H, alkyl; R5 = H, alkyl, cyano; R6 = H, alkyl, aryl, heterocyclyl; R7-R11 = H, OH, alkyl, alkoxy, aryl, cycloalkyl, heterocyclyl], pharmaceutically acceptable salts, solvates or hydrate thereof, were prepared for use as antitumor agents. Thus, epothilone derivative

II was prepared from 2,3-dehydro epothilone A, via silylation of hydroxyl group, potassium cyanide addition, followed by deprotection. The prepared epothilone derivs. were assayed *in vitro* for their effect on tubulin polymerization and for cytotoxicity against HCT-116 human colon

carcinoma cells. Therapeutic compns. containing I or in combination with other therapeutic agents useful in the treatment of cancer or other proliferative diseases are also claimed.

IT 476623-89-9P 476623-90-2P 476623-91-3P
476623-92-4P


RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of epothilone derivs. for therapeutic use as anticancer agents)

RN 476623-89-9 HCAPLUS

CN 4,17-Dioxabicyclo[4.1.0]heptadecane-7-carbonitrile,
11-hydroxy-8,8,10,12-tetramethyl-3-[(1E)-1-methyl-2-(2-methyl-4-
thiazolyl)ethenyl]-5,9-dioxo-, (1S,3S,7S,10R,11S,12S,16R)- (CA INDEX
NAME)


Absolute stereochemistry.
Double bond geometry as shown.



RN 476623-90-2 HCPLUS

CN 4,17-Dioxabicyclo[14.1.0]heptadecane-7-carbonitrile,
 11-hydroxy-8,8,10,12-tetramethyl-3-[(1E)-1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-5,9-dioxo-, (1S,3S,7R,10R,11S,12S,16R)- (CA INDEX NAME)


Absolute stereochemistry.
 Double bond geometry as shown.



RN 476623-91-3 HCPLUS

CN 4,17-Dioxabicyclo[14.1.0]heptadecane-7-carbonitrile,
 11-hydroxy-8,8,10,12,16-pentamethyl-3-[(1E)-1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-5,9-dioxo-, (1S,3S,7S,10R,11S,12S,16R)- (CA INDEX NAME)


Absolute stereochemistry.
 Double bond geometry as shown.



RN 476623-92-4 HCAPLUS

CN 4,17-Dioxabicyclo[14.1.0]heptadecane-7-carbonitrile,
 11-hydroxy-8,8,10,12,16-pentamethyl-3-[(1E)-1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-5,9-dioxo-, (1S,3S,7R,10R,11S,12S,16R)- (CA INDEX NAME)

Absolute stereochemistry.
 Double bond geometry as shown.



OS.CITING REF COUNT: 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD
(2 CITINGS)

REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L9 ANSWER 9 OF 16 HCPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2003:396890 HCPLUS

DOCUMENT NUMBER: 138:385210

TITLE: Preparation of novel epothilone derivatives via bioconversion for use in pharmaceutical compositions for the treatment of cancer and non-cancer hyperproliferative disorders

INVENTOR(S): Tang, Li; Metcalf, Brian; Katz, Leonard; Ashley, Gary W.; Lau, Janice; Licari, Peter

PATENT ASSIGNEE(S): Kosan Biosciences, Inc., USA

SOURCE: PCT Int. Appl., 67 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003042217	A2	20030522	WO 2002-US36814	20021114 <--
WO 2003042217	A3	20031002		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW			

RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG

AU 2002363803 A1 20030526 AU 2002-363803 20021114 <--
US 20030219877 A1 20031127 US 2002-295342 20021114 <--
US 7070964 B2 20060704

PRIORITY APPLN. INFO.: US 2001-334734P P 20011115
WO 2002-US36814 W 20021114

GI



AB Epothilone derivs., such as I (R = H, Me; R1 = F, OH), were prepared via a combination of bioconversion and synthetic methods for therapeutic uses, such as the treatment of cancer, non-cancer hyperproliferative disorders, multiple sclerosis, rheumatoid arthritis, atherosclerosis, and restenosis. Organisms used for the bioformation of epothilones include *Streptomyces hygroscopicus* ATCC 55098, *Amycolata autotrophica* ATCC 35203, *Actinomyces* sp. strain SC15847, *Saccharopolyspora erythrea* NRRL2338, *Saccharopolyspora erythrea* K39-14, *Myxococcus xanthus* K 111-72, *Myxococcus xanthus* K111-76 and *Myxococcus xanthus* K111-78. Thus, epothilone D, as well as its hydroxylated derivs., were prepared by a fermentation process using *Saccharopolyspora erythrea* K39-14. Other epothilone derivs., such as 11(R)- and 11(S)-fluoroepothilones, were prepared from the hydroxylated derivs. using synthetic reaction schemes. Pharmaceutical compns. and dosages of the prepared epothilones were presented.

IT 152044-53-6P, Epothilone A 152044-54-7P, Epothilone

B


RL: BPN (Biosynthetic preparation); PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of novel epothilone derivs. via bioconversion for use in pharmaceutical compns. for treatment of cancer and non-cancer hyperproliferative disorders)

RN 152044-53-6 HCPLUS

CN 4,17-Dioxabicyclo[4.1.0]heptadecane-5,9-dione, 7,11-dihydroxy-8,10,12-tetramethyl-3-[(1E)-1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-, (1S,3S,7S,10R,11S,12S,16R)- (CA INDEX NAME)


Absolute stereochemistry. Rotation (-).
Double bond geometry as shown.



RN 152044-54-7 HCPLUS

CN 4,17-Dioxabicyclo[4.1.0]heptadecane-5,9-dione,
7,11-dihydroxy-8,10,12,16-pentamethyl-3-[(1E)-1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-, (1S,3S,7S,10R,11S,12S,16R)- (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).
 Double bond geometry as shown.



OS.CITING REF COUNT: 3 THERE ARE 3 CAPLUS RECORDS THAT CITE THIS RECORD

(3 CITINGS)

REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L9 ANSWER 10 OF 16 HCPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2003:77585 HCPLUS

DOCUMENT NUMBER: 138:137091

TITLE: Preparation of epothilone derivatives for therapeutic use as antitumor agents

INVENTOR(S): Ashley, Gary; Metcalf, Brian

PATENT ASSIGNEE(S): USA

SOURCE: U.S. Pat. Appl. Publ., 38 pp.

CODEN: USXXCO

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1



PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20030023082	A1	20030130	US 2002-145405	20020513 <--
PRIORITY APPLN. INFO.:			US 2001-291242P	P 20010515
			US 2001-309099P	P 20010731

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OTHER SOURCE(S): MARPAT 138:137091

GI



AB The present invention relates to the preparation of epothilone deriva., such as I and II [R = H, OH, NH2; R8 = H, C1-5-aliphatic;

R10 = H, Cl-5-aliphatic, aryl; A-B = CHCH, C:C; D = O, S, NR10, CR10:N, CONH, etc.; W = O, NR8; Y = H, heterocyclyl, for pharmaceutical use as antiproliferative and antitumor agents. These epothilone derivs. can be used for the treatment of diseases or conditions characterized by undesired cellular hyperproliferation, such as cancer, atrophic gastritis, inflammatory hemolytic anemia, graft rejection, inflammatory neutropenia, bullous pemphigoid, coeliac disease, demyelinating neuropathies, dermatomyositis, inflammatory bowel disease, multiple sclerosis, myocarditis, myositis, nasal polyps, chronic sinusitis, pemphigus vulgaris, primary glomerulonephritis, psoriasis, surgical adhesions, stenosis or restenosis, scleritis, scleroderma, eczema, periodontal disease, polycystic kidney disease, and type I diabetes. Thus, 26-(imidazol-2-yl)Epothilone D II (R = H, W = O, Y = 2-imidazolyl) by treating 30,70-bis(trimethylsilyl)- derivative of 26-(methoxymethylene)epothilone D II (R = H, W = O, Y = :CHOMe) with glyoxal and ammonium acetate in THF. I.v. and liposomal pharmaceutical formulations and a pretreatment regimen for Cremophor toxicity were presented.

IT 371979-65-6P, 9-Oxoepothilone B


RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(preparation of epothilone derivs. for therapeutic use as antitumor and antiproliferative agents)

RN 371979-65-6 HCAPLUS

CN 4,17-Dioxabicyclo[14.1.0]heptadecane-5,9,13-trione,
7,11-dihydroxy-8,8,10,12,16-pentamethyl-3-[(1E)-1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-, (1S,3S,7S,10R,11S,12R,16R)- (CA INDEX NAME)

Absolute stereochemistry.
Double bond geometry as shown.




OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD
(1 CITINGS)

L9 ANSWER 11 OF 16 HCPLUS COPYRIGHT 2010 ACS on STN
 ACCESSION NUMBER: 2001:886112 HCPLUS
 DOCUMENT NUMBER: 136:5855
 TITLE: Preparation of epothilone
 derivatives for pharmaceutical use
 in the treatment of cancer and other disorders
 characterized by cellular hyperproliferation
 INVENTOR(S): Santi, Daniel; Fardis, Maria; Ashley, Gary
 PATENT ASSIGNEE(S): Kosan Biosciences, Inc., USA
 SOURCE: PCT Int. Appl., 87 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001092255	A2	20011206	WO 2001-US15763	20010515 <--
WO 2001092255	A3	20020228		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
US 20020045609	A1	20020418	US 2001-859085	20010515 <--
PRIORITY APPLN. INFO.:			US 2000-207655P	P 20000526
			US 2000-218260P	P 20000714
			US 2000-231552P	P 20000911

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OTHER SOURCE(S): MARPAT 136:5855
 GI



AB Epothilone derivs., such as I [R = Me, CH₂OH, CHO; R4 = H, OH, oxo, amino, etc.; R5 = H, OH, oxo, R6 = H, OH, oxo, alkyl, alkylester, halogen, etc.; R7 = H, alkyl, halogen, hydroxyalkyl, alkoxyalkyl, arylalkyl, heterocyclylalkyl, etc.; R8 = R9 = H; R8R9 = bond, O; R5R6 = bond, W = O, NR11; R11 = H, alkyl, aryl], were prepared for therapeutic use in the treatment of cancer and non-cancer disorders characterized by cellular hyperproliferation. Thus, (11S)-hydroxyepothilone D II (R6 = α -OH) and its (11R)-diastereomer II (R = β -OH) were prepared by hydroxylation of epothilone D using SeO₂ and Me₃COOH by stirring in CH₂Cl₂ for 48 h. The prepared epothilone derivs. were assayed for cytotoxicity against MCF-7 breast, MDR breast, SF-268 glioma and NCI-H460 lung cancer cell lines and were assayed for tubulin polymerization inhibition.

IT 371979-65-6P


RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(preparation and formulation of epothilone derivs. for pharmaceutical use in the treatment of cancer and other disorders characterized by cellular hyperproliferation)

RN 371979-65-6 HCPLUS

CN 4,17-Dioxabicyclo[14.1.0]heptadecane-5,9,13-trione,
7,11-dihydroxy-8,10,12,16-pentamethyl-3-[(1E)-1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-, (1S,3S,7S,10R,11S,12R,16R)- (CA INDEX NAME)

Absolute stereochemistry.
Double bond geometry as shown.



OS.CITING REF COUNT: 12 THERE ARE 12 CAPLUS RECORDS THAT CITE THIS RECORD (12 CITINGS)
 REFERENCE COUNT: 8 THERE ARE 8 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L9 ANSWER 12 OF 16 HCAPLUS COPYRIGHT 2010 ACS on STN
 ACCESSION NUMBER: 2001:780370 HCAPLUS
 DOCUMENT NUMBER: 135:331294
 TITLE: Preparation of epothilone derivatives for pharmaceutical use in the treatment of cancer
 INVENTOR(S): Buchmann, Bernd; Klar, Ulrich; Skuballa, Werner; Schwede, Wolfgang; Hoffmann, Jens; Lichtner, Rosemarie
 PATENT ASSIGNEE(S): Schering A.-G., Germany
 SOURCE: Ger. Offen., 42 pp.
 CODEN: GWXXBX
 DOCUMENT TYPE: Patent
 LANGUAGE: German
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 10020517	A1	20011025	DE 2000-10020517	20000419 <--
WO 2001081342	A2	20011101	WO 2001-EP4552	20010419 <--
WO 2001081342	A3	20020510		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW				

RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG

EP 1276740 A2 20030122 EP 2001-936262 20010419 <--

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR

JP 2003531207 T 20031021 JP 2001-578432 20010419 <--



NO 2002005029 A 20021018 NO 2002-5029 20021018 <--

US 20040058969 A1 20040325 US 2002-257925 20021018 <--

PRIORITY APPLN. INFO.: DE 2000-10020517 A 20000419
WO 2001-EP4552 W 20010419

OTHER SOURCE(S): MARPAT 135:331294

GI




AB Epothilones, such as I [R3 = heteroaryl, heteroarylalkenyl, heteroarylhaloalkenyl, etc.; R8, R8a = H, alkyl, arylalkyl; R8R8a = alkylene, heteroalkene; R10 = H, alkyl, alkynyl, alkynyl; R1R16a = bond, O; R16 = H, CN, alkyl, halogen; X = O, NH; X1 = O, CH2], were prepared for a variety of therapeutic uses, such as treatment of malignant tumors, proliferative diseases, leukemia, and chronic inflammatory diseases. Thus, epothilone II was prepared via a multistep synthetic sequence starting from (3S)-dihydro-3-hydroxy-4,4-dimethyl-2(3H)-furanone, L-(-)-malic acid, and [(2-methyl-4-thiazolyl)methyl]phosphonic acid di-Et ester. Pharmaceutical formulations of the prepared oxa-epothilones were discussed, but specific biol. activity data was not presented.

IT 369646-19-5P

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
(preparation of epothilone derivs. for pharmaceutical use in the treatment of cancer)

RN 369646-19-5 HCPLUS
 CN 4,17-Dioxabicyclo[4.1.0]heptadecane-16-carbonitrile,
 7,11-dihydroxy-8,8,10,12-tetramethyl-3-[(1E)-1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-5,9-dioxo-, (3S,7S,10R,11S,12S)- (CA INDEX NAME)

Absolute stereochemistry.
 Double bond geometry as shown.



OS.CITING REF COUNT: 5 THERE ARE 5 CAPLUS RECORDS THAT CITE THIS RECORD
 (5 CITINGS)

L9 ANSWER 13 OF 16 HCPLUS COPYRIGHT 2010 ACS on STN
 ACCESSION NUMBER: 2000:790507 HCPLUS
 DOCUMENT NUMBER: 133:362656
 TITLE: Preparation of 6-alkenyl-, 6-alkynyl- and
 6-epoxyepothilone derivatives and their antitumor
 activity
 INVENTOR(S): Klar, Ulrich; Schwede, Wolfgang; Skuballa, Werner;
 Buchmann, Bernd; Hoffmann, Jens; Lichtner, Rosemarie
 Schering Aktiengesellschaft, Germany
 PATENT ASSIGNEE(S): PCT Int. Appl., 298 pp.
 SOURCE: CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 3
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2000066589	A1	20001109	WO 2000-IB657	20000501 <--
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU,				

LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE,
 SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW
 RW: GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE,
 DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF,
 CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG
 DE 19921086 A1 20001102 DE 1999-19921086 19990430 <--
 DE 19954228 A1 20010913 DE 1999-19954228 19991104 <--
 DE 10015836 A1 20011011 DE 2000-10015836 20000327 <--
 CA 2371226 A1 20001109 CA 2000-2371226 20000501 <--
 BR 2000010190 A 20020108 BR 2000-10190 20000501 <--
 EP 1173441 A1 20020123 EP 2000-922826 20000501 <--
 EP 1173441 B1 20090826
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, MC, PT, IE,
 SI, LT, LV, FI, RO, CY, SE
 JP 2002543203 T 20021217 JP 2000-615619 20000501 <--
 JP 4024003 B2 20071219
 EE 200100568 A 20030217 EE 2001-568 20000501 <--
 NZ 514989 A 20040227 NZ 2000-514989 20000501 <--
 AU 772750 B2 20040506 AU 2000-43103 20000501 <--
 SK 286858 B6 20090605 SK 2001-1551 20000501
 AT 440847 T 20090915 AT 2000-922826 20000501
 IN 2001MN01305 A 20070504 IN 2001-MN1305 20011019
 BG 106053 A 20020531 BG 2001-106053 20011026 <--
 BG 65601 B1 20090227
 NO 2001005278 A 20011221 NO 2001-5278 20011029 <--
 MX 2001011039 A 20030630 MX 2001-11039 20011030 <--
 US 7125893 B1 20061024 US 2002-979939 20020606 <--
 HK 1046681 A1 20080829 HK 2002-108204 20021113
 IN 2005MN00837 A 20070608 IN 2005-MN837 20050802
 US 20060046997 A1 20060302 US 2005-214988 20050831 <--
 PRIORITY APPLN. INFO.:
 DE 1999-19921086 A1 19990430
 DE 1999-19954228 A1 19991104
 DE 2000-10015836 A1 20000327
 DE 2000-10013363 A 20000309
 WO 2000-IB657 W 20000501
 IN 2001-MN1305 A3 20011019
 US 2002-979939 A3 20020606

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OTHER SOURCE(S): MARPAT 133:362656

GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB The antitumor agents, 6-alkenyl-, 6-alkynyl- and 6-epoxyepothilones I
 (R1a, R1b are same or different = H, C1-C10 alkyl, C6-C12 aryl, C7-C20
 aralkyl each optionally substituted; or together = $(CH_2)^m$ m = 1-5 or
 $-CH_2OCH_2-$; R2a(R2b replace with b) = H, substituted alkyl, aryl,
 aralkyl, $(CH_2)^ra-C_6H_4-C_6H_4-$ or $=C_6H_4-C_6H_4-C_6H_4-$ pa-R26a, Q, Q1 where n = 0-5; ra,
 rb = the same or different and = 0-4; pa, pb = the same or different and =
 0-3; R3a = H, substituted alkyl, aryl or aralkyl; R3b = OH, OPG14; R14 =
 H, OR14a, halogen and R14a = H, SO2-alkyl, SO2-aryl or SO2-aralkyl; R4 =
 H, substituted alkyl, aryl or aralkyl, halogen, OR25, CN; R26a, R26b =
 same or different = H, substituted alkyl, aryl or aralkyl, C1-C10 acyl or
 if pa or pb > 0, addnl. a group OR27; R25 = R27 = R22 = H, PG; R5 = H,

substituted alkyl, aryl or aralkyl, $(\text{CH}_2)_s\text{T}$ $s = 1-4$, $\text{T} = \text{OR22}$ or halogen; $\text{R}6$, $\text{R}7 = \text{H}$ or together = bond or O ; $\text{G} = \text{X}=\text{CR}8$ or bi- or tricyclic aryl radical and $\text{R}8 = \text{H}$, halogen, CN , or substituted alkyl, aryl or aralkyl; $\text{X} = \text{O}$, two OR23 groups, $\text{C}_2\text{-C}_{10}$ -alkylene- α,ω -dioxy straight chain or branched; $\text{H/CR}9$ or $\text{CR}10\text{R}11$ group and $\text{R}23 =$ alkyl radical, $\text{R}9 = \text{H}$, PG , $\text{R}10, \text{R}11 =$ same or different = H , substituted alkyl, aryl or aralkyl, or together with the methylene are a 5-7 carbocyclic ring; $\text{D-E} = \text{CH}_2\text{CH}_2$ or OCH_2 ; $\text{A} = \text{OC}(\text{O})$, OCH_2 , $\text{CH}_2\text{C}(\text{O})$, $\text{NR}29\text{C}(\text{O})$, $\text{NR}29\text{SO}_2$ and $\text{R}29 = \text{H}$, alkyl; $\text{Z} = \text{O}$ or $\text{H/CR}12$ and $\text{R}12 = \text{H}$, PG) were prepared. Thus II was prepared in a multistep synthesis starting from $(4S)-4-(2\text{-methyl-1-oxoprop-2-yl})-2,2\text{-dimethyl}[1,3]\text{dioxane}$ and 5-trimethylsilylpent-4-in-1-yl magnesium bromide. II had an IC_{50} value [nM] of 3.0 for the growth inhibition of human MCF-7 breast- and 75 for multidrug resistant NCI/ADR carcinoma cell lines with a selectivity of 2.5. The new epothilone derivs.



interact with tubulin by stabilizing microtubuli that are formed. They are able to influence the cell-splitting in a phase-specific manner and are therefore useful in treating diseases or conditions associated with the need for cell growth, division and/or proliferation. Thus the epothilone derivs. are suitable for treating malignant tumors, e.g., ovarian, stomach, colon, adeno-, breast, lung, head and neck carcinomas, malignant melanoma, acute lymphocytic and myelocytic leukemia; and for anti-angiogenesis therapy as well as for treatment of chronic inflammatory diseases (such as psoriasis, arthritis).

IT 305840-25-9P 305840-26-0P 305840-30-6P
305840-31-7P

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
(preparation of 6-alkenyl-, 6-alkynyl- and 6-epoxyepothilone derivs. and their use in pharmaceutical preps.)



RN 305840-25-9 HCPLUS
CN 4,17-Dioxabicyclo[14.1.0]heptadecane-5,9-dione,
7,11-dihydroxy-8,8,12,16-tetramethyl-3-[(1E)-1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-10-(2-propen-1-yl)-, (1S,3S,7S,10R,11S,12S,16R)- (CA INDEX NAME)

Absolute stereochemistry.
Double bond geometry as shown.



RN 305840-26-0 HCAPLUS
 CN 4,17-Dioxabicyclo[4.1.0]heptadecane-5,9-dione,
 7,11-dihydroxy-8,12,16-tetramethyl-3-(1E)-1-methyl-2-(2-methyl-4-
 thiazolyl)ethenyl-10-(2-propen-1-yl)-, (1R,3S,7S,10R,11S,12S,16S)- (CA
 INDEX NAME)



Absolute stereochemistry.
 Double bond geometry as shown.



RN 305840-30-6 HCAPLUS

CN 4,17-Dioxabicyclo[14.1.0]heptadecane-5,9-dione,
 7,11-dihydroxy-8,8,12,16-tetramethyl-3-(1E)-1-methyl-2-(2-methyl-4-thiazolyl)ethenyl-10-(2-propen-1-yl)-, (1R,3S,7S,10R,11S,12S,16R)- (CA
 INDEX NAME)



Absolute stereochemistry.
 Double bond geometry as shown.



RN 305840-31-7 HCAPLUS

CN 4,17-Dioxabicyclo[4.1.0]heptadecane-5,9-dione,
 7,11-dihydroxy-8,8,12,16-tetramethyl-3-(1E)-1-methyl-2-(2-methyl-4-thiazolyl)ethenyl-10-(2-propen-1-yl)-, (1S,3S,7S,10R,11S,12S,16S)- (CA
 INDEX NAME)

Absolute stereochemistry.
 Double bond geometry as shown.



OS.CITING REF COUNT: 9 THERE ARE 9 CAPLUS RECORDS THAT CITE THIS RECORD
 (9 CITINGS)
 REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS
 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L9 ANSWER 14 OF 16 HCAPLUS COPYRIGHT 2010 ACS on STN
 ACCESSION NUMBER: 1999:12688 HCAPLUS
 DOCUMENT NUMBER: 130:196529
 TITLE: Preparation of new epothilone
 derivatives as pharmaceutical agents
 INVENTOR(S): Klar, Ulrich; Schwede, Wolfgang; Skuballa, Werner;
 Buchmann, Bernd; Schirner, Michael
 PATENT ASSIGNEE(S): Schering Aktiengesellschaft, Germany
 SOURCE: PCT Int. Appl., 185 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: German
 FAMILY ACC. NUM. COUNT: 4
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
------------	------	------	-----------------	------

WO 9907692	A2	19990218	WO 1998-EP5064	19980810 <--
WO 9907692	A3	19990514		
W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DK, EE, ES, FI, GB, GE, GH, GM, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW				
RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
DE 19735574	A1	19990211	DE 1997-19735574	19970809 <--
DE 19735575	A1	19990211	DE 1997-19735575	19970809 <--
DE 19735578	A1	19990211	DE 1997-19735578	19970809 <--
DE 19748928	A1	19990429	DE 1997-19748928	19971024 <--
DE 19749717	A1	19990506	DE 1997-19749717	19971031 <--
DE 19751200	A1	19990520	DE 1997-19751200	19971113 <--
DE 19813821	A1	19990923	DE 1998-19813821	19980320 <--
CA 2299608	A1	19990218	CA 1998-2299608	19980810 <--
AU 9893409	A	19990301	AU 1998-93409	19980810 <--
EP 1005465	A2	20000607	EP 1998-946309	19980810 <--
EP 1005465	B1	20070725		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, CY, AL, MK				
JP 2001512723	T	20010828	JP 2000-506196	19980810 <--
ZA 9810403	A	20000515	ZA 1998-10403	19981113 <--
IN 190805	A1	20030823	IN 1998-DE3413	19981116 <--
US 20030144523	A1	20030731	US 2000-485292	20000503 <--
US 7407975	B2	20080805		
IN 2002DE01305	A	20050311	IN 2002-DE1305	20021227
US 20090018342	A1	20090115	US 2008-178039	20080723 <--
PRIORITY APPLN. INFO.:			DE 1997-19735574	A 19970809
			DE 1997-19735575	A 19970809
			DE 1997-19735578	A 19970809
			DE 1997-19748928	A 19971024
			DE 1997-19749717	A 19971031
			DE 1997-19751200	A 19971113
			DE 1998-19813821	A 19980320
			WO 1998-EP5064	W 19980810
			US 2000-485292	A1 20000503

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OTHER SOURCE(S): MARPAT 130:196529

GI



AB Epothilone derivs. of formula I [X = O, alkylene- α , ω -dioxy, two alkoxy groups, etc.; Y = O, H₂; Z = O, (H, OH), (H, protected OH); R1a, R1b = H, alkyl, aryl, aralkyl, or together = (CH₂)_m where m = 2, 3, 4, 5; R2a, R2b = H, alkyl, aryl, aralkyl, or together = (CH₂)_n where n = 2, 3, 4, 5; when D-E = CH₂CH₂ or when Y = O, R2a or R2b may not be H/Me; R3 = H, alkyl, aryl, aralkyl; R4a, R4b = H, alkyl, aryl, aralkyl, or together = (CH₂)_p where p = 2, 3, 4, 5; D-E = CH₂CH₂, CH₂CH, C.tplbond.C, 2,3-oxiranediyl, CH(OH)CH(OH), CH(OH)CH₂; R5 = H, alkyl, aryl, aralkyl; R6, R7 = H, together = a saturated bond or O; R8 = H, alkyl, aryl, aralkyl all of which may be substituted] are prepared. Thus, the title compds. (4S,7R,8S,9S,13E,16S(E))- and (4S,7R,8S,9S,13Z,16S(E))-4,8-dihydroxy-7-ethyl-16-(1-methyl-2-(2-methyl-4-thiazolyl)ethenyl)-1-oxa-5,9,13-tetramethylcyclohexadec-13-en-2,6-dione (II) were prepared in many steps. The new compds. interact with tubulin by stabilizing formed microtubuli. They are capable of influencing cell division in a phase-specific manner and are suitable for the treatment of malignant tumors, such as ovarian, gastric, colon, breast, lung, head and neck carcinoma, adenocarcinoma, malignant melanoma, and acute lymphocytic and myelocytic leukemia. They are also suited for anti-angiogenesis therapy and for the treatment of chronic inflammatory diseases (psoriasis, arthritis). To prevent uncontrolled cell growth on, and for better tolerability of, medical implants, the derivs. can be introduced into or applied to polymeric materials. The compds. provided for in the invention can be used alone or, to achieve additive or synergistic effects, in combination with other principles and substance categories used in tumor therapy.

IT 220773-47-7P


RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); RCT (Reactant); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); RACT (Reactant or reagent); USES (Uses)

(preparation of epothilone derivs. as antitumor agents)

RN 220773-47-7 HCAPLUS

CN 4,17-Dioxabicyclo[14.1.0]heptadecane-5,9-dione,
10-ethyl-7,11-dihydroxy-8,8,12,16-tetramethyl-3-[(1E)-1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-, (1S,3S,7S,10R,11S,12S,16R)- (CA INDEX NAME)

Absolute stereochemistry.
Double bond geometry as shown.



IT 220773-48-8P 220773-49-9P 220773-50-2P
 220773-61-5P 220773-64-8P 220773-65-9P
 220773-66-0P 220773-67-1P 220776-29-4P
 220776-30-7P 220776-31-8P 220776-32-9P
 220776-44-3P 220776-45-4P 220776-46-5P
 220776-47-6P 220776-50-1P 220776-51-2P
 220776-52-3P 220776-53-4P


RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of epothilone derivs. as antitumor agents)

RN 220773-48-8 HCPLUS

CN 4,17-Dioxabicyclo[14.1.0]heptadecane-5,9-dione,
 10-ethyl-7,11-dihydroxy-8,8,12,16-tetramethyl-3-[(1E)-1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-, (1R,3S,7S,10R,11S,12S,16S)- (CA INDEX NAME)

Absolute stereochemistry.


Double bond geometry as shown.



RN 220773-49-9 HCPLUS


CN 4,17-Dioxabicyclo[14.1.0]heptadecane-5,9-dione,
 10-ethyl-7,11-dihydroxy-8,8,12,16-tetramethyl-3-[(1E)-1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-, (1R,3S,7S,10R,11S,12S,16R)- (CA INDEX NAME)

Absolute stereochemistry.
 Double bond geometry as shown.




RN 220773-50-2 HCAPLUS
 CN 4,17-Dioxabicyclo[4.1.0]heptadecane-5,9-dione,
 10-ethyl-7,11-dihydroxy-8,8,12,16-tetramethyl-3-[(1E)-1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-, (1S,3S,7S,10R,11S,12S,16S)- (CA INDEX NAME)

Absolute stereochemistry.
 Double bond geometry as shown.



RN 220773-61-5 HCAPLUS
 CN 4,17-Dioxabicyclo[4.1.0]heptadecane-5,9-dione,
 10-ethyl-7,11-dihydroxy-8,8,12,16-tetramethyl-3-[(1E)-1-methyl-2-(2-methyl-3-oxido-4-thiazolyl)ethenyl]-, (1S,3S,7S,10R,11S,12S,16R)- (CA INDEX NAME)


Absolute stereochemistry.
 Double bond geometry as shown.



RN 220773-64-8 HCAPLUS

CN 4,17-Dioxabicyclo[14.1.0]heptadecane-5,9-dione,
 7,11-dihydroxy-8,8,12,16-tetramethyl-3-[(1E)-1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-10-(phenylmethyl)-, (1S,3S,7S,10R,11S,12S,16R)- (CA INDEX NAME)


Absolute stereochemistry.
 Double bond geometry as shown.



RN 220773-65-9 HCAPLUS

CN 4,17-Dioxabicyclo[14.1.0]heptadecane-5,9-dione,
 7,11-dihydroxy-8,12,16-tetramethyl-3-[(1E)-1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-10-(phenylmethyl)-, (1R,3S,7S,10R,11S,12S,16S)- (CA
 INDEX NAME)


Absolute stereochemistry.
 Double bond geometry as shown.



RN 220773-66-0 HCAPLUS

CN 4,17-Dioxabicyclo[14.1.0]heptadecane-5,9-dione,
 7,11-dihydroxy-8,8,12,16-tetramethyl-3-[1E)-1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-10-(phenylmethyl)-, (1S,3S,7S,10R,11S,12S,16S)- (CA
 INDEX NAME)


Absolute stereochemistry.
 Double bond geometry as shown.



RN 220773-67-1 HCAPLUS

CN 4,17-Dioxabicyclo[14.1.0]heptadecane-5,9-dione,
 7,11-dihydroxy-8,8,12,16-tetramethyl-3-[1(E)-1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-10-(phenylmethyl)-, (1R,3S,7S,10R,11S,12S,16R)- (CA
 INDEX NAME)


Absolute stereochemistry.
 Double bond geometry as shown.



RN 220776-29-4 HCAPLUS

CN 4,17-Dioxabicyclo[14.1.0]heptadecane-5,9-dione,
7,11-dihydroxy-8,8,10,16-tetramethyl-3-[(1E)-1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-12-(trifluoromethyl)-, (1S,3S,7S,10R,11R,12S,16R)- (CA
INDEX NAME)


Absolute stereochemistry.
Double bond geometry as shown.



RN 220776-30-7 HCAPLUS

CN 4,17-Dioxabicyclo[14.1.0]heptadecane-5,9-dione,
 7,11-dihydroxy-8,8,10,16-tetramethyl-3-[(1E)-1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-12-(trifluoromethyl)-, (1R,3S,7S,10R,11R,12S,16R)- (CA INDEX NAME)


Absolute stereochemistry.
 Double bond geometry as shown.



RN 220776-31-8 HCAPLUS

CN 4,17-Dioxabicyclo[14.1.0]heptadecane-5,9-dione,
 7,11-dihydroxy-8,8,10,16-tetramethyl-3-[(1E)-1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-12-(trifluoromethyl)-, (1R,3S,7S,10R,11R,12S,16S)- (CA INDEX NAME)


Absolute stereochemistry.
 Double bond geometry as shown.



RN 220776-32-9 HCAPLUS

CN 4,17-Dioxabicyclo[14.1.0]heptadecane-5,9-dione,
 7,11-dihydroxy-8,8,10,16-tetramethyl-3-[(1E)-1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-12-(trifluoromethyl)-, (1S,3S,7S,10R,11R,12S,16S)- (CA
 INDEX NAME)


Absolute stereochemistry.
 Double bond geometry as shown.



RN 220776-44-3 HCAPLUS

CN 4,17-Dioxabicyclo[14.1.0]heptadecane-5,9-dione,
 7,11-dihydroxy-8,10,12-tetramethyl-3-[(1E)-1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-16-(trifluoromethyl)-, (1S,3S,7S,10R,11S,12S,16R)- (CA INDEX NAME)


Absolute stereochemistry.
 Double bond geometry as shown.



RN 220776-45-4 HCAPLUS

CN 4,17-Dioxabicyclo[14.1.0]heptadecane-5,9-dione,
 7,11-dihydroxy-8,10,12-tetramethyl-3-[(1E)-1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-16-(trifluoromethyl)-, (1R,3S,7S,10R,11S,12S,16R)- (CA INDEX NAME)


Absolute stereochemistry.
 Double bond geometry as shown.



RN 220776-46-5 HCAPLUS

CN 4,17-Dioxabicyclo[14.1.0]heptadecane-5,9-dione,
7,11-dihydroxy-8,8,10,12-tetramethyl-3-[(1E)-1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-16-(trifluoromethyl)-, (1R,3S,7S,10R,11S,12S,16S)- (CA INDEX NAME)


Absolute stereochemistry.
Double bond geometry as shown.



RN 220776-47-6 HCAPLUS

CN 4,17-Dioxabicyclo[4.1.0]heptadecane-5,9-dione,
 7,11-dihydroxy-8,8,10,12-tetramethyl-3-[(1E)-1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-16-(trifluoromethyl)-, (1S,3S,7S,10R,11S,12S,16S)- (CA INDEX NAME)

Absolute stereochemistry.
 Double bond geometry as shown.





RN 220776-50-1 HCAPLUS

CN 4,17-Dioxabicyclo[4.1.0]heptadecane-5,9-dione,
7,11-dihydroxy-8,10,12-tetramethyl-3-[(1E)-1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-16-(1,1,2,2,2-pentafluoroethyl)-,
(IS,3S,7S,10R,11S,12S,16R)- (CA INDEX NAME)

Absolute stereochemistry.
Double bond geometry as shown.

PAGE 1-A







RN 220776-51-2 HCPLUS

CN 4,17-Dioxabicyclo[14.1.0]heptadecane-5,9-dione,
 7,11-dihydroxy-8,8,10,12-tetramethyl-3-[(1E)-1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-16-(1,1,2,2,2-pentafluoroethyl)-,
 (1R,3S,7S,10R,11S,12S,16R)- (CA INDEX NAME)

Absolute stereochemistry.

Double bond geometry as shown.




RN 220776-52-3 HCPLUS

CN 4,17-Dioxabicyclo[4.1.0]heptadecane-5,9-dione,
 7,11-dihydroxy-8,8,10,12-tetramethyl-3-(1E)-1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-16-(1,1,2,2,2-pentafluoroethyl)-,


(1R, 3S, 7S, 10R, 11S, 12S, 16S)- (CA INDEX NAME)

Absolute stereochemistry.
Double bond geometry as shown.

PAGE 1-A



PAGE 1-B




10591921

RN 220776-53-4 HCPLUS
CN 4,17-Dioxabicyclo[14.1.0]heptadecane-5,9-dione,
7,11-dihydroxy-8,8,10,12-tetramethyl-3-[(1E)-1-methyl-2-(2-methyl-4-
thiazolyl)ethenyl]-16-(1,1,2,2,2-pentafluoroethyl)-,
(1S,3S,7S,10R,11S,12S,16S)- (CA INDEX NAME)

Absolute stereochemistry.
Double bond geometry as shown.

PAGE 1-A





OS.CITING REF COUNT: 22 THERE ARE 22 CAPLUS RECORDS THAT CITE THIS RECORD (22 CITINGS)
 REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L9 ANSWER 15 OF 16 HCAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 1997:456769 HCAPLUS

DOCUMENT NUMBER: 127:50474

ORIGINAL REFERENCE NO.: 127:9629a

TITLE: Preparation of epothilone derivatives as agrochemicals and pharmaceuticals

INVENTOR(S): Hoefie, Gerhard; Kiffe, Michael

PATENT ASSIGNEE(S): Gesellschaft fuer Biotechnologische Forschung Mbh (Gbf), Germany

SOURCE: Ger. Offen., 9 pp.

CODEN: GWXXBX

DOCUMENT TYPE: Patent

LANGUAGE: German

FAMILY ACC. NUM. COUNT: 2

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 19542986	A1	19970522	DE 1995-19542986	19951117 <--
WO 9719086	A1	19970529	WO 1996-EP5080	19961118 <--
W: JP, US				
RW: AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
EP 873341	A1	19981028	EP 1996-939097	19961118 <--
EP 873341	B1	20030910		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,				

IE, FI


EP 903348	A1	19990324	EP 1998-121523	19961118 <--
EP 903348	B1	20020605		
EP 903348	B2	20080827		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,				
IE, FI				
JP 2000500757	T	20000125	JP 1997-519381	19961118 <--
JP 4183099	B2	20081119		
EP 1186606	A1	20020313	EP 2001-127352	19961118 <--
EP 1186606	B1	20040317		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,				
IE, FI				
AT 218556	T	20020615	AT 1998-121523	19961118 <--
PT 903348	E	20021129	PT 1998-121523	19961118 <--
ES 2178093	T3	20021216	ES 1998-121523	19961118 <--
AT 249463	T	20030915	AT 1996-939097	19961118 <--
PT 873341	E	20040227	PT 1996-939097	19961118 <--
AT 261961	T	20040415	AT 2001-127352	19961118 <--
ES 2206607	T3	20040516	ES 1996-939097	19961118 <--
EP 1440973	A2	20040728	EP 2004-5011	19961118 <--
EP 1440973	A3	20041020		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,				
IE, FI				
PT 1186606	E	20040831	PT 2001-127352	19961118 <--
ES 2218328	T3	20041116	ES 2001-127352	19961118 <--
US 6288237	B1	20010911	US 1998-77055	19980803 <--
US 20010034452	A1	20011025	US 2001-836134	20010416 <--
US 6613912	B2	20030902		
US 20040087634	A1	20040506	US 2003-602770	20030625 <--
US 6831076	B2	20041214		
DE 1995-19542986 A 19951117				
DE 1996-19639456 A 19960925				
EP 1996-939097 A3 19961118				
EP 2001-127352 A3 19961118				
WO 1996-EP5080 W 19961118				
US 1998-77055 A3 19980803				
US 2001-836134 A3 20010416				

PRIORITY APPLN. INFO.:


ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OTHER SOURCE(S): MARPAT 127:50474

GI




Absolute stereochemistry. Rotation (-).
Double bond geometry as shown.



RN 152044-54-7 HCAPLUS

14,17-Dioxabicyclo[4.1.0]heptadecane-5,9-dione,
7,11-dihydroxy-8,10,12,16-pentamethyl-3-[(1E)-1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-, (15,3S,7S,10R,11S,12S,16R)- (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).
Double bond geometry as shown.




OS.CITING REF COUNT: 9 THERE ARE 9 CAPLUS RECORDS THAT CITE THIS RECORD
(12 CITINGS)

L9 ANSWER 16 OF 16 HCAPLUS COPYRIGHT 2010 ACS on STN
 ACCESSION NUMBER: 1997:443365 HCAPLUS
 DOCUMENT NUMBER: 127:81289
 ORIGINAL REFERENCE NO.: 127:15595a,15588a
 TITLE: Preparation of epothilone
 derivatives as agrochemicals and
 pharmaceuticals
 INVENTOR(S): Hofle, Gerhard; Kiffe, Michael
 PATENT ASSIGNEE(S): Gesellschaft Fur Biotechnologische Forschung Mbh
 (Gbf), Germany; Hofle, Gerhard; Kiffe, Michael
 SOURCE: PCT Int. Appl., 38 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: German
 FAMILY ACC. NUM. COUNT: 2
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9719086	A1	19970529	WO 1996-EP5080	19961118 <--
W: JP, US RW: AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
DE 19542986	A1	19970522	DE 1995-19542986	19951117 <--
DE 19639456	A1	19980326	DE 1996-19639456	19960925 <--
EP 873341	A1	19981028	EP 1996-939097	19961118 <--
EP 873341	B1	20030910		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				
JP 2000500757	T	20000125	JP 1997-519381	19961118 <--
JP 4183099	B2	20081119		
AT 249463	T	20030915	AT 1996-939097	19961118 <--
US 6288237	B1	20010911	US 1998-77055	19980803 <--
US 20040087634	A1	20040506	US 2003-602770	20030625 <--
US 6831076	B2	20041214		
PRIORITY APPLN. INFO.:				
			DE 1995-19542986 A 19951117	
			DE 1996-19639456 A 19960925	
			WO 1996-EP5080 W 19961118	
			US 1998-77055 A3 19980803	
			US 2001-836134 A3 20010416	

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OTHER SOURCE(S): MARPAT 127:81289
GI



AB The title compds., e.g., I [R = H, C1-4 alkyl; R1, R2 = H, C1-6 alkyl, C1-6 acyl, benzoyl, C1-4 trialkylsilyl, benzyl, Ph, C1-6 alkoxy, C6 alkyl-, hydroxy-, and halo-substituted benzyl or phenyl; X, Y = H, halo, pseudohalo, OH, acyloxy, alkoxy, benzoyloxy; or YZ = O, bond; however, I may not be epothilone A or B], useful as agrochems. and pharmaceuticals (no data), are prepared. Thus, epothilone A in acetone containing trifluoroacetic acid was heated overnight at 50° and the reaction mixture was adjusted to pH 7 with 1 M phosphate buffer to give 2 isomers, each in 19% yield.


IT 152044-53-6, Epothilone A 152044-54-7, Epothilone B

RL: RCT (Reactant); RACT (Reactant or reagent)
(preparation of epothilone derivs. as agrochems. and pharmaceuticals)


RN 152044-53-6 HCPLUS

CN 4,17-Dioxabicyclo[14.1.0]heptadecane-5,9-dione,
7,11-dihydroxy-8,8,10,12-tetramethyl-3-[1(E)-1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-, (1S,3S,7S,10R,11S,12S,16R)- (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).
Double bond geometry as shown.



RN 152044-54-7 HCPLUS

CN 4,17-Dioxabicyclo[14.1.0]heptadecane-5,9-dione,
7,11-dihydroxy-8,8,10,12,16-pentamethyl-3-[(1E)-1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-, (1S,3S,7S,10R,11S,12S,16R) - (CA INDEX NAME)Absolute stereochemistry. Rotation (-).
Double bond geometry as shown.

10591921

OS.CITING REF COUNT: 26 THERE ARE 26 CAPLUS RECORDS THAT CITE THIS
RECORD (30 CITINGS)
REFERENCE COUNT: 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> log y
COST IN U.S. DOLLARS SINCE FILE TOTAL
ENTRY SESSION
FULL ESTIMATED COST 122.06 313.82
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS) SINCE FILE TOTAL
ENTRY SESSION
CA SUBSCRIBER PRICE -13.60 -13.60

STN INTERNATIONAL LOGOFF AT 15:27:35 ON 04 JAN 2010